



Computational analysis of deposition and translocation of inhaled nicotine and acrolein in the human body with e-cigarette puffing topographies

Ahmadreza Haghnegahdar, Yu Feng, Xiaole Chen & Jiang Lin

To cite this article: Ahmadreza Haghnegahdar, Yu Feng, Xiaole Chen & Jiang Lin (2018) Computational analysis of deposition and translocation of inhaled nicotine and acrolein in the human body with e-cigarette puffing topographies, *Aerosol Science and Technology*, 52:5, 483-493, DOI: [10.1080/02786826.2018.1447644](https://doi.org/10.1080/02786826.2018.1447644)

To link to this article: <https://doi.org/10.1080/02786826.2018.1447644>

 View supplementary material 

 Accepted author version posted online: 02 Mar 2018.
Published online: 26 Mar 2018.

 Submit your article to this journal 

 Article views: 151

 View related articles 

 View Crossmark data 



Computational analysis of deposition and translocation of inhaled nicotine and acrolein in the human body with e-cigarette puffing topographies

Ahmadreza Haghnegahdar^a, Yu Feng^a, Xiaole Chen^b, and Jiang Lin^c

^aSchool of Chemical Engineering, Oklahoma State University, Stillwater, Oklahoma, USA; ^bSchool of Energy and Environment, Southeast University, Nanjing, Jiangsu, China; ^cCollege of Light Industry, Zhejiang University of Science and Technology, Hangzhou, Zhejiang, China

ABSTRACT

Recently, toxicants such as formaldehyde and acrolein were detected in electronic cigarette (EC) aerosols. It is imperative to conduct research and provide sufficient quantitative evidence to address the associated potential health risks. However, it is still a lack of informative data, i.e., high-resolution local dosimetry of inhaled aerosols in lung airways and other systemic regions, due to the limited imaging resolutions, restricted operational flexibilities, and invasive nature of experimental and clinical studies. In this study, an experimentally validated multiscale numerical model, i.e., Computational Fluid-Particle Dynamics (CFPD) model combined with a Physiologically Based Toxicokinetic (PBTK) model is developed to predict the systemic translocation of nicotine and acrolein in the human body after the deposition in the respiratory system. *In-silico* parametric analysis is performed for puff topography influence on the deposition and translocation of nicotine and acrolein in human respiratory systems and the systemic region. Results indicate that the puff volume and holding time can contribute to the variations of the nicotine and acrolein plasma concentration due to enhanced aerosol deposition in the lung. The change in the holding time has resulted in significant difference in the chemical translocation which was neglected in a large group of experimental studies. The capability of simulating multiple puffs of the new CFPD-PBTK model paves the way to a valuable computational simulation tool for assessing the chronic health effects of inhaled EC toxicants.

ARTICLE HISTORY

Received 3 October 2017
Accepted 25 February 2018

EDITOR

Yifang Zhu

1. Introduction

Electronic cigarettes (ECs), perceived as the safer alternatives to conventional cigarettes, are manufactured in a wide variety of device designs with widely varying nicotine concentrations, added flavors, and glycerol (VG), as well as propylene glycol (PG) to imitate smoke via vaporization instead of combustion (Allen et al. 2016). However, studies also claimed that some e-cigarette products have immediate adverse physiological effects after short-term use, e.g., an increase in impedance, peripheral airway flow resistance, and oxidative stress among healthy smokers who used an e-cigarette for 5 min (Alfi and Talbot 2013), while the long-term health effects are still unknown and worthy of further investigations. Recently, toxicants such as formaldehyde, acetaldehyde, acrolein, diacetyl, benzaldehyde, and vanillin were also detected in e-cigarette aerosols (Goniewicz, Hajek, & McRobbie 2014; Allen et al. 2016; Flora et al. 2017). These toxicants are either generated by the low-level

thermal degradation or exist in flavorings. Because of the potential health impact of the newly discovered chemical compounds listed above (Alarcon 1976; De Woskin et al. 2003; Wang et al. 2008; Mishra et al. 2015; Morris et al. 2015; Abraham et al. 2011), it is imperative to conduct research and provide sufficient quantitative evidence to address the potential health risks for the optimization of e-cigarette product design or facilitate regulatory approval. Thus, it is critical to define reliable biomarkers and understand how those chemical compounds transport, translocate and accumulate in the human body. However, it is still a lack of informative data, i.e., high-resolution local dosimetry of inhaled aerosols in lung airways and other systemic regions. The knowledge and data gap are mainly due to the limited imaging resolutions, restricted operational flexibilities, and invasive nature of experimental and clinical studies. Indeed, since ECs have not been in widespread use sufficiently long, the human exposure studies until now are limited to

CONTACT Yu Feng yu.feng@okstate.edu School of Chemical Engineering, Oklahoma State University, 420 Engineering North, Stillwater, OK 74078, USA.
Color versions of one or more of the figures in the article can be found online at www.tandfonline.com/uast.

Supplemental data for this article can be accessed on the [publisher's website](#).

very short-term health responses. Meanwhile, although animal studies were performed to investigate the health effects on mice and rats due to the exposure to EC aerosols, observations are very restrictive regarding imaging resolution and operational flexibility (Morris 2012).

As an alternative, *in-silico* study using high-fidelity numerical methods, i.e., Computational Fluid-Particle Dynamics (CFPD) plus Physiologically Based Toxicokinetic (PBTk) model (Figure 1), are capable of providing valuable dosimetry data of multicomponent EC aerosols in subject-specific respiratory tracts (Chen et al. 2017; Haghnegahdar and Feng 2017; Kleinstreuer and Feng 2013). Furthermore, a high-quality CFPD-PBTk model is also a promising whole-body dosimetry prediction tool, which is noninvasive, cost-effective, and time-saving compared to *in-vitro* and *in-vivo* investigations. Compared to those lumped parameter approaches and semi-empirical models, the CFPD-PBTk model is developed based on the natural law of physics and chemistry with fewer assumptions and simplifications. Thus, an advanced noninvasive, reliable, and generalized engineering tool based on the multiscale CFPD-PBTk

framework is developed and employed in this study to analyze the comparative health risk reductions of EC aerosol components, i.e., nicotine and acrolein, with different puffing topographies. This *in-silico* study has relevance to public health because it will enable a major health question to be answered by simulations in a short period, i.e., what is the relationship between the chronic passive e-cigarette vaping of adolescents and their health risks. The integration of multiple disciplinary types of research will bridge the gap of EC aerosol exposure and cardiopulmonary health effect assessments, leading ultimately to a better understanding of the complexity and dynamics of EC aerosol transport, deposition and translocation in human bodies.

Efforts have been made towards the development of numerical models to predict the fates of inhaled tobacco products. The necessary approaches to have a reliable model that can represent the essential governing physics are presented in the review by Kleinstreuer and Feng (2013). Simulations of the EC generated aerosol with the multipath particle deposition model (MMPD) approach developed by Manigrasso et al. (2015). The connection




Figure 1. Framework of the CFPD-PBTk model.

of the CFPD and the PBTK demands additional description that can present the rule of the interconnecting tissue layer. Kimbell et al. (1993) has completed the single-phase formaldehyde transport in Fisher rat nasal cavity. In this work, the complete water solubility of the formaldehyde led them to consider the zero concentration condition at the airway walls, and consequently, the diffusion process was uncoupled from those in the surrounding tissues. Deposition and mucosa layer translocation of vapor in the nasal cavity with hybrid CFD-PBTK simulations were studied by Morris, et al. (1993). Bush et al. (1998) simulated non-reactive vapors uptake in the rat nasal cavity by including compartmental designs for respired air, the mucus layer, the epithelial cell layer, and the sub-epithelial layer containing blood vessels. Corley et al. (2012, 2015) have reported the connected CFD and PBTK model for a single vapor phase deposition of the cigarette constituent in the respiratory system of the three systems of rat, monkey, and human considering mucus, epithelial, and submucosal layers with diffusion and associated reactions.

In this paper, the new multiscale CFPD-PBTK model is developed and employed to predict the local fates of nicotine and acrolein from entering the human upper airways to the systemic regions. Since EC aerosol composition generated varies due to different puffing behaviors and EC products (Lisko et al. 2015). Hence, the parametric analysis of puff pattern influence on transport and translocation dynamics is performed for individualized health risk assessments. EC Puffing topographies were selected as inlet conditions based on existing investigations and standards (Cobb et al. 2010; Goniewicz et al. 2014; Goniewicz et al. 2017; Shahab et al. 2017). Simulations of multiple puffs with holdings are enabled and performed for the first time using the CFPD-PBTK model.

The layout of this article is as follows: Section 2 introduces the framework of the multiscale CFPD-PBTK model; Section 3 contains the numerical results of multi-component EC aerosol transport, deposition, and translocation with insightful discussion and parametric analysis. Section 4 summarizes the findings of the present study. Section 5 proposes future work to improve the generalized feasibility of the CFPD-PBTK modeling framework. Numerical setup, mesh independence tests, model validations, and the limitations of this study are provided as the online supplementary information (SI).

2. Methodology

Governed by the conservation laws of nature, the Computational Fluid-Particle Dynamics (CFPD) and

Physiologically Based Toxicokinetic (PBTK) model is a promising tool to assess the chronic exposure risks of EC aerosols and provide informative and high-resolution data promptly. The schematic of the CFPD-PBTK modeling framework is shown in Figure 1. Integrating the multiscale model validations and optimizations, the CFPD-PBTK model provides local information about how different levels of puffing may affect the deposition and translocation of toxicants in both lung and systemic regions. The CFPD model is developed based on Euler-Lagrange scheme (Feng et al. 2016) specifically for multi-component EC aerosol dynamics in an idealized human upper airway model from mouth to Generation 3 (G3). The existence of the dominant chemicals of the nicotine, acrolein, formaldehyde, vegetable glycerin (VG), and propylene glycol (PG) are tracked both in the particle and vapor forms. Also, the wall of the respiratory system is considered as a sink with fractional to complete absorption for the uptake of the chemicals into the systemic region. The PBTK model for inhaled toxicants is developed and validated. It is assumed that the toxicant's distribution through blood flow with the biological structure of tissues are homogeneous rate-limited diffusion (Robinson et al. 1992). The important mechanisms including absorption, distribution, metabolism, and excretion in each organ for each toxicant are considered. Physiologic parameters (cardiac output, ventilation rate, blood flow rate to the organs and organ volumes) are obtained and optimized accordingly. The system of governing equations and boundary conditions are provided in the SI.

3. Results and discussion

3.1. EC Particle transport and deposition

The translocation of the EC toxicants including acrolein and nicotine are numerically investigated in the idealized human upper airway model. Three realistic puffing inlet conditions are applied (see SI for details). As shown in Figures 2–7, the local EC particle deposition patterns are significantly influenced by airflow regime determined by the average puff flow rate. Moreover, after the puffing duration, the holding time will result in increased deposition due to Brownian motion and the gravity sedimentation effects. The transient local airflow patterns with secondary flows and recirculation regions are shown in Figures 2 and 3. Specifically, Figures 2a–j show the airflow pattern transition at the sagittal plane and frontal plane by visualizing velocity magnitudes at different time steps. It can be observed from Figures 2a–d that a jet flow is evolved from the




Figure 2. Transient airflow patterns in the idealized human upper airway model ($Q_{in} = 55$ ml per 3.0 s): (a) $t = 0.05$ s; (b) $t = 0.10$ s; (c) $t = 0.20$ s; (d) $t = 0.30$ s; (e) $t = 0.40$ s; (f) $t = 0.50$ s; (g) $t = 0.75$ s; (h) $t = 1.00$ s; (i) $t = 2.00$ s; (j) $t = 3.00$ s.

EC mouthpiece at the beginning of the puff. At $t = 0.3$ s, the high-velocity jet impacts the lower palate forms the recirculation region in the oral cavity, and deviate the path towards pharynx and trachea. At $t = 0.40$ s the inlet jet is completely formed in the oral cavity and the pharynx (Figure 2e) and the second recirculation region at the front side of the pharynx is formed due to the centrifugal effects of the mainstream flow. At $t = 0.50$ s, the high-velocity jet reaches the trachea region. The third recirculation is formed at the back of the trachea due to the pressure variations induced by the glottis. Due to the small particle size, most particles

will follow the high-velocity jet and have small chance to enter the recirculation regions. However, there is a small portion of the particles, which can enter the recirculation region. Those particles will have longer residence time and higher chance to deposit induced by Brownian motion effects. It should be noted that the backflow will increase the residence time of particles entering the recirculating regions, while reducing the local deposition avoid direct impaction of the particles carried by the mainstream flow (Figure 3).

Additionally, Figures 3a–e show the transient local deposition patterns of EC particles. The deposition of




Figure 3. Local deposition patterns of EC particles ($Q_{in} = 55$ ml per 3.0 s with 2-s holding): (a) $t = 1.0$ s; (b) $t = 2.0$ s; (c) $t = 3.0$ s; (d) $t = 4.0$ s; (e) $t = 5.0$ s.

the particles at $t = 1.0$ s, 2.0 s, and 3.0 s followed by 1.0 s and 2.0 s holdings are provided. Low deposition efficiencies (DEs) at the airway walls covered by recirculation regions. Enhanced local deposition efficiencies (LDEs) are found at the lower palate, pharynx, and bifurcating points because of inertial impaction and gravitational sedimentation. Figures 3d and e demonstrate that particles continue to deposit at lung airways during the holding phase. Increased holding time will result in higher depositions. Therefore, it is necessary to simulate

the holding duration to obtain accurate particle deposition patterns for EC vaping and passive exposure.

3.2. EC vapor species transport and absorption

As an example, Figures 4 and 5 show the mass fraction contour evolution of acrolein and nicotine vapors at the sagittal plane for puff volume of 55 ml. After being inhaled, acrolein and nicotine vapors follow the airflow well to the tracheobronchial




Figure 4. Mass fraction distribution of acrolein vapor at different ($Q_{in} = 55$ ml per 3.0 s): (a) $t = 0.05$ s; (b) $t = 0.10$ s; (c) $t = 0.20$ s; (d) $t = 0.30$ s; (e) $t = 0.40$ s; (f) $t = 0.50$ s; (g) $t = 0.75$ s; (h) $t = 1.00$ s; (i) $t = 2.00$ s; (j) $t = 3.00$ s.




Figure 5. Mass fraction distribution of nicotine vapor at different ($Q_{in} = 55$ ml per 3.0 s): (a) $t = 0.05$ s; (b) $t = 0.10$ s; (c) $t = 0.20$ s; (d) $t = 0.30$ s; (e) $t = 0.40$ s; (f) $t = 0.50$ s; (g) $t = 0.75$ s; (h) $t = 1.00$ s; (i) $t = 2.00$ s; (j) $t = 3.00$ s.

region gradually. Due to the high diffusion rates of both vapors into the airway tissues, most of the acrolein (99%) and nicotine (99%) in vapor forms are absorbed before reaching the glottis (see SI for the binary diffusivities). Therefore, the translocation of both vapor species into the blood circulation happens at human upper airways.

3.3. Puff topography influence on EC aerosol transport dynamics

The impact of puff topography on EC aerosol transport and deposition were investigated and shown in Figures 6 and 7. As shown in Figure 6 for two puff volumes, most particles deposit on the mouth to glottis region for both inlet puffing volumes. Higher puffing flow rate induce longer residence time of particles in G1 to G3, leading to higher deposition due to inertial impaction. “Hot spots” of particle deposition include lower palate, pharynx, glottis, and bifurcating points of each generation.

Additionally, comparisons of regional deposition fractions (RDFs) for all cases with different inlet conditions are shown in Figure 7. The histogram indicates that high puff flow rate leads to enhanced RDFs due to the stronger inertial impaction effect and less EC particles will be able to transport into deeper airways. Increased holding time will also increase RDFs because of the increased

particle residence time. The subsequent holdings to active puffs increase the residence time of the EC particles, which were still suspending in the lung. Hence, the regional deposition will continuously occur with the progress of holding time.

3.4. Translocations of EC toxicants in the systemic region

As the lung input to the PBTK model, total toxicant deposition was calculated by adding depositions of both particulate and vapor phases. Optimized values of bio-availability factors were determined for the simplified CFPD-PBTK connection model, by matching plasma concentration profiles from benchmark experiments (Ramôa et al. 2015; Shahab et al. 2017; Goniewicz et al. 2017). Since CFPD results indicate that the existence of the acrolein and nicotine in the region after trachea is negligible (see Section 3.2), it can be concluded that the total uptake of both nicotine and acrolein are mostly dependent on their absorption in human upper airways. Distribution and translocation of the toxicants are discussed separately in the following sections.

3.4.1. Nicotine translocation

The concentration input profile in both vapor phase and particulate phase for nicotine are provided for all




Figure 6. Comparison of the EC particle deposition patterns at $t = 3.0$ s between two puffing volumes.

simulation cases with different puff topographies, i.e., two puffing volumes (55 and 80 ml per 3.0 s) and three holding times (0.0 s, 2.0 s, and 7.0 s). Figures 8a and b show the concentration profiles for 10 s including 3 s puffing time followed by 2 and 7 s holding time for nicotine at two puffing volumes of 55 and 80 ml. The AUC over 30-s puffing duration for each case is also provided which shows the average uptake value of each puff. Particle depositions are scaled up based on the realistic particle number concentrations. Subsequently, nicotine input

concentration due to particle deposition and vapor absorption are considered, and the correlation is generated using nonlinear regression by the least square method.

The AUCs show that for the case of 55 ml puffing volume the average uptake concentration per puff can be significantly higher for 2.0 s holding time (67.13% increase) and for 7 s holding time (112.49% increase). Also for the case with 80 ml puffing volume, the increases are 57.52% and 95.57%, respectively, for 2 and 7 s holding times. Moreover, increasing puff volume will increase vapor absorption. However, it has little influence on particle deposition. As a result, the total deposition of the nicotine and, respectively, its uptake has increased by 21.61% for the case without holding time. This difference between different puffing volumes is simulated as 14.62% and 11.93% for 2 and 7 s holding times respectively in the course of 30.0 s puff duration. In this study, the influence of the exhalation has not been considered, and the deposition concentration after the holding time is set to zero. This assumption is reasonable since most deposition of nicotine in both vapor and particle forms occur before $t = 10.0$ s (Figures 8a and b).

The predicted nicotine plasma concentrations under different puff volumes are provided in Figures 9a and b. 10 puffs with 30.0 s puffing duration are considered. The first puff results in a sudden increase of nicotine concentration. The subsequent 9 puffs increase the concentration gradually with a slower rate. The peak plasma concentrations after 10 puffs for the case of 55 and 80 ml puffing volumes are 15.58 ng/ml and 18.88 ng/ml without considering holding time. It can also be observed that due to the higher absorption and deposition, the increase in holding time generates higher plasma concentrations. At the same time, the realistic holding time will result in a change of



Figure 7. Comparisons of EC particle deposition fractions in the idealized human upper airway model.





Figure 8. Time courses of nicotine uptake concentration after 3-s puffing duration followed by 7 s holding time: (a) 55 ml puffing volume and (b) 80 ml puffing volume for 10 s of the total puffing duration (30 s).

the exhalation time. Overall, the increase in holding times will contribute to the exponential increase of the nicotine bioavailability. Changing puffing volume only has a slight effect on nicotine plasma concentration compared to changing holding time.

3.4.2. Acrolein translocation

Acrolein exists only in the vapor phase based on the initial condition applied in this study. As a result, the uptake concentration will be induced by the vapor phase absorption, which is shown in Figure 10 for the two cases with 55 and 80 ml puff volumes. It is obvious that the increase in puff volume results in the increase of the peak value of the uptake concentration from 0.508 to





Figure 9. Time courses of nicotine plasma concentration: (a) 55 ml puffing volume and (b) 80 ml puffing volume (the nicotine uptake concentration has been scaled down for better visualization).

0.792 ng/ml. The average AUC for 30 s for different cases with different holding times are also provided. For the cases with of 55 ml and 80 ml puffing volumes, increases of AUC as representative of per puff exposure after considering two holding times of 2.0 s and 7.0 s are 45.51%, 83.27%, 44.45%, and 134.53, respectively.

Figures 11a and b present the acrolein plasma concentration within 10 puffs. The puff by puff increase of the acrolein concentration can be observed. The first puff has resulted in major increase in the concentration. After 10 puffs, peak concentrations of acrolein with different holding times from 0.0 s to 7.0 s are 0.001221, 0.001648, 0.001803 ng/ml for 55 ml puffing volume case and 0.001947, 0.002614, 0.002864 ng/ml for 80 ml puffing




Figure 10. Time course of acrolein uptake concentration after 3-s puffing duration followed by 7 s holding time at two puffing volumes of 55 and 80 ml for 10 s of the total puffing duration (30 s).

volume case. For better visualization effect, the acrolein uptakes were scaled down by 4×10^{-3} for better visualization. The reason for the difference is that the difference of the considered lymph vein volume compared to the plasma volume.

The formation of 3-HPMA produced by the binding between acrolein and glutathione has been identified in the urinary sample of rats (Alarcon 1976). As a result, 3-HPMA is considered as the main metabolite and possible biomarker of acrolein bioavailability. 3-HPMA concentration at the kidney is shown in Figure 12. It should be noted that in this study the regional bioavailability factors are considered as representative of reactions taking place to form the metabolites of the species. In this way, complete conversion of acrolein to 3-HPMA has been initiated to the PBTK model for the acrolein. It is worth mentioning that acrolein itself has lipophilic characteristic and the perfusion-limited models can be reliable prediction based on our simulation. However, it is also important for the future studies to investigate the fate of transport in the tissue for its main metabolites. In Figure 12, the puff-by-puff increase of the concentration in the kidney can be observed. Damped concentration profile in the kidney can be attributed to the configuration of the compartments. The peak concentration in excretion for 55 ml puffing volume are 110.74, 160.40, and 201.58 ng/mg creatinine for the three cases with 0.0, 2.0, and 7.0 s holdings after the 3.0 s puff duration. Also, for cases with 80 ml puffing volume, the peak values are 176.78, 245.35, and 321.15 ng/mg creatinine, respectively.

Based on the simulation the excretion profile shows a peak value that is suggested by the experiments (Shahab et al. 2017). The increase in puff volume from 55 to 80 ml for a case without holding time has contributed to 59.63%





Figure 11. Time courses of acrolein plasma concentration: (a) 55 ml puffing volume and (b) 80 ml puffing volume (the acrolein uptake concentration has been scaled down for better visualization).




Figure 12. Time courses of the ratio between excreted 3-HPMA and creatinine in the kidney with different puff topographies.

increase. However, by increasing the holding time for the case with 55 ml puffing volume, the peak value increase is approximately 82.03%. This difference can depict the importance of the holding time that can contribute to significant change in translocation magnitude in the human body. Existing standards of the puffing machines have focused mostly on three parameters: puff volume, duration and interval time. Therefore, the importance of the additional parameter that must also be investigated in the future experiments. Other parameters may include EC product type (McNeill et al. 2015), and interaction between EC chemicals (Kane and Alarie 1978).

4. Conclusions

In this study, a multiscale CFPD-PBTK model is developed and applied to accurately predict multicomponent EC aerosol transport, deposition, and translocation from the human respiratory system to systemic regions. Specifically, aerosol dynamics was modeled by the CFPD model established based on the Euler-Lagrange scheme. A 9-compartment PBTK framework subsequently simulates the translocation of nicotine and acrolein in the systemic region. Employing the CFPD-PBTK model, it is the first time that *in-silico* investigations can reflect realistic EC vaping scenarios, i.e., multiple puffs with real EC user behaviors. The model advances the field of aerosol exposure science to pave the way to a valuable computational simulation tool for assessing the long-term health effects of inhaled e-cigarette toxicants in the human respiratory pathway and systemic regions. Quantitative data of EC toxicant fates to the health endpoints are generated and discussed. Based on the numerical results, conclusions and novel insight are summarized as follows:

- (1) Most of the vaporized nicotine and acrolein are absorbed in the upper airway from mouth to G1. In contrast, EC aerosol particle deposition occurs in all regions from mouth to G3.
- (2) Both particulate and vapor forms of nicotine and acrolein contribute to the deposition and translocation in the human body.
- (3) The increase from 55 to 80 ml puffing volume per 3 s results in significant increase of the peak nicotine and acrolein plasma concentration as 21.2% and 59.45% for the cases without holding.
- (4) With fixed puffing volume, increasing holding time will result in the increase of peak plasma concentration for both nicotine and acrolein.
- (5) Holding time between consecutive puffs has significant impacts on the EC aerosol deposition and translocation. Therefore, it is necessary to include holding time in both experimental and numerical studies to provide accurate predictions.

5. Future work

Future work will focus on the improvement of the CFPD-PBTK model considering more complex aerosol dynamics. The next generation CFPD-PBTK model will include the coagulation, condensation, and evaporation among phases. The interconnection model will be developed for different pulmonary regions, to reflect the different diffusion and reaction characteristics. Additionally, subject-specific human airway configurations will be used for vulnerable subpopulation groups to EC products, e.g., adolescents, and COPD patients. To improve the current PBTK model, the couplings among compartments will be considered to simulate the realistic time delay effects in different organs. Also, a diffusion-limited PBTK model will be developed for formaldehyde and other toxicants in EC aerosols.

Funding

Research reported in this publication was supported by the National Institute of General Medical Sciences of the National Institute of Health under Award Number P20GM103648. The authors also gratefully acknowledge the financial support of Dr. Jiang Lin from the National Natural Science Foundation of China (No. 51246002). The use of ANSYS software (Canonsburg, PA, USA) as part of the ANSYS-OSU academic partnership agreement is gratefully acknowledged (Dr. Thierry Marchal, Global Industry Director). Some of the computing for this project was performed at the OSU High Performance Computing Center at Oklahoma State University (Dr. Dana Brunson, Director and Dr. Evan Linde, Research Cyberinfrastructure Analyst).

References

- Abraham, K., Andres, S., Palavinkas, R., Berg, K., Appel, K. E., & Lampen, A. (2011). Toxicology and risk assessment of acrolein in food. *Mol. Nutr Food Res.*, 55(9):1277–1290. doi:10.1002/mnfr.201100481.
- Alarcon, R. (1976). Studies on the *in vivo* formation of acrolein: 3-hydroxy-propylmercapturic acid as an index of cyclophosphamide (NSC-26271) activation. *Cancer Treat Rep.*, 60(4):327–335.
- Alfi, M., & Talbot, P. (2013). Health-related effects reported by electronic cigarette users in online forums. *J Med Internet Res.*, 15(4).
- Allen, J. G., Flanigan, S. S., LeBlanc, M., Vallarino, J., MacNaughton, P., Stewart, J. H., & Christiani, D. C. (2016). Flavoring chemicals in e-cigarettes: diacetyl, 2, 3-pentanedione, and acetoin in a sample of 51 products, including fruit-, candy-, and cocktail-flavored e-cigarettes. *Environ Health Perspect.*, 124(6):733. doi:10.1289/EHP348.
- Andersen, M., Sarangapani, R., Gentry, R., Clewell, H., Covington, T., & Frederick, C. B. (2000). Application of a hybrid CFD-PBPK nasal dosimetry model in an inhalation risk assessment: an example with acrylic acid. *Toxicol Sci.*, 57(2):312–325. doi:10.1093/toxsci/57.2.312.

- Belka, M., Lizal, F., Jedelsky, J., Jicha, M., & Pospisil, J. (2017). *Measurement of an electronic cigarette aerosol size distribution during a puff*. Paper presented at the EPJ Web of Conferences.
- Bush, M. L., Frederick, C. B., Kimbell, J. S., & Ultman, J. S. (1998). A CFD-PBPK hybrid model for simulating gas and vapor uptake in the rat nose. *Toxicol Applied Pharmacol.*, 150(1):133–145. doi:10.1006/taap.1998.8407.
- Chen, X., Feng, Y., Zhong, W., & Kleinstreuer, C. (2017). Numerical investigation of the interaction, transport and deposition of multicomponent droplets in a simple mouth-throat model. *J Aerosol Sci.*, 105:108–127. doi:10.1016/j.jaerosci.2016.12.001.
- Cobb, N. K., Byron, M. J., Abrams, D. B., & Shields, P. G. (2010). Novel nicotine delivery systems and public health: the rise of the “e-cigarette”. In: *American Public Health Association*.
- Corley, R. A., Kabilan, S., Kuprat, A. P., Carson, J. P., Jacob, R. E., Minard, K. R., ... Glenny, R. (2015). Comparative risks of aldehyde constituents in cigarette smoke using transient computational fluid dynamics/physiologically based pharmacokinetic models of the rat and human respiratory tracts. *Toxicol Sci.*, 146(1):65–88. doi:10.1093/toxsci/kfv071.
- Corley, R. A., Kabilan, S., Kuprat, A. P., Carson, J. P., Minard, K. R., Jacob, R. E., ... Cox, T. (2012). Comparative computational modeling of airflows and vapor dosimetry in the respiratory tracts of rat, monkey, and human. *Toxicol Sci.*, 128(2):500–516. doi:10.1093/toxsci/kfs168.
- De Woskin, R., Greenberg, M., Pepelko, W., & Strickland, J. (2003). Toxicological review of acrolein (cas no. 107-02-08) in support of summary information on the integrated risk information system (Iris). *Washington, DC: US Environmental Protection Agency*.
- Feng, Y., Kleinstreuer, C., Castro, N., & Rostami, A. (2016). Computational transport, phase change and deposition analysis of inhaled multicomponent droplet-vapor mixtures in an idealized human upper lung model. *J Aerosol Sci.*, 96:96–123. doi:10.1016/j.jaerosci.2016.03.001.
- Flora, J. W., Wilkinson, C. T., Wilkinson, J. W., Lipowicz, P. J., Skapars, J. A., Anderson, A., & Miller, J. H. (2017). Method for the determination of carbonyl compounds in e-cigarette aerosols. *Journal of Chromatographic Science*, 55(2):142–148.
- Goniewicz, M. L., Gawron, M., Smith, D. M., Peng, M., Jacob, P., & Benowitz, N. L. (2017). Exposure to nicotine and selected toxicants in cigarette smokers who switched to electronic cigarettes: a longitudinal within-subjects observational study. *Nicotine Tob Res.*, 19(2):160–167. doi:10.1093/ntr/ntw160.
- Goniewicz, M. L., Hajek, P., & McRobbie, H. (2014). Nicotine content of electronic cigarettes, its release in vapour and its consistency across batches: regulatory implications. *Addiction*, 109(3):500–507. doi:10.1111/add.12410.
- Haghnegahdar, A., Feng, Y. (2017). The translocation of nicotine from human lung to systemic regions due to e-cigarette aerosol inhalation: a numerical study. *5th International Conference on Computational and Mathematical Biomedical Engineering (CMBE)*, Pittsburgh, PA, USA.
- Ingebretsen, B. J., Cole, S. K., & Alderman, S. L. (2012). Electronic cigarette aerosol particle size distribution measurements. *Inhal Toxicol.*, 24(14):976–984. doi:10.3109/08958378.2012.744781.
- Kane, L. E., & Alarie, Y. (1978). Evaluation of sensory irritation from acrolein-formaldehyde mixtures. *Am Indus Hygiene Assoc J.*, 39(4):270–274. doi:10.1080/0002889778507758.
- Kimbell, J., Gross, E., Joyner, D., Godo, M., & Morgan, K. (1993). Application of computational fluid dynamics to regional dosimetry of inhaled chemicals in the upper respiratory tract of the rat. *Toxicol Applied Pharmacol.*, 121(2):253–263. doi:10.1006/taap.1993.1152.
- Kleinstreuer, C., & Feng, Y. (2013). Lung deposition analyses of inhaled toxic aerosols in conventional and less harmful cigarette smoke: a review. *Int J Environ Res Public Health.*, 10(9):4454–4485. doi:10.3390/ijerph10094454.
- Lisko, J. G., Tran, H., Stanfill, S. B., Blount, B. C., & Watson, C. H. (2015). Chemical composition and evaluation of nicotine, tobacco alkaloids, pH, and selected flavors in e-cigarette cartridges and refill solutions. *Nicotine Tob Res.*, 17(10):1270–1278. doi:10.1093/ntr/ntu279.
- Manigrasso, M., Buonanno, G., Fuoco, F. C., Stabile, L., & Avino, P. (2015). Aerosol deposition doses in the human respiratory tree of electronic cigarette smokers. *Environ Pollut.*, 196:257–267. doi:10.1016/j.envpol.2014.10.013.
- McNeill, A., Brose, L. S., Calder, R., Hitchman, S. C., Hajek, P., & McRobbie, H. (2015). E-cigarettes: an evidence update.
- Mishra, A., Chaturvedi, P., Datta, S., Sinukumar, S., Joshi, P., & Garg, A. (2015). Harmful effects of nicotine. *Ind J Med Paediat Oncol: official journal of Indian Society of Medical & Paediatric Oncology*, 36(1):24. doi:10.4103/0971-5851.151771.
- Morris, J. B. (2012). Biologically-based modeling insights in inhaled vapor absorption and dosimetry. *Pharmacology & therapeutics*, 136(3):401–413.
- Morris, J. B., Hassett, D. N., & Blanchard, K. T. (1993). A physiologically based pharmacokinetic model for nasal uptake and metabolism of nonreactive vapors. *Toxicol Applied Pharmacol.*, 123(1):120–129. doi:10.1006/taap.1993.1228.
- Morris, P. B., Ference, B. A., Jahangir, E., Feldman, D. N., Ryan, J. J., Bahrami, H., ... Al-Mallah, M. H. (2015). Cardiovascular effects of exposure to cigarette smoke and electronic cigarettes. *J Am College Cardiol.*, 66(12):1378–1391. doi:10.1016/j.jacc.2015.07.037.
- Ramôa, C. P., Hiler, M. M., Spindle, T. R., Lopez, A. A., Kar-aoghlanian, N., Lipato, T., ... Eissenberg, T. (2015). Electronic cigarette nicotine delivery can exceed that of combustible cigarettes: a preliminary report. *Tob control, tobaccocontrol-2015-052447*.
- Robinson, D. E., Balter, N. J., & Schwartz, S. L. (1992). A physiologically based pharmacokinetic model for nicotine and cotinine in man. *J Pharmacokinet Biopharma.*, 20(6):591–609. doi:10.1007/BF01064421.
- Shahab, L., Goniewicz, M. L., Blount, B. C., Brown, J., McNeill, A., Alwis, K. U., ... West, R. (2017). Nicotine, Carcinogen, and Toxin Exposure in Long-Term E-Cigarette and Nicotine Replacement Therapy Users: A Cross-sectional Study. *E-Cigarettes and Toxin Exposure. Ann Internal Med.*, 166(6):390–400. doi:10.7326/M16-1107.
- Smith, J., Van Ness, H., & Abbott, M. (1996). Chemical engineering thermodynamics. *Sat.*, 18:1–3.
- Uchiyama, S., Ohta, K., Inaba, Y., & Kunugita, N. (2013). Determination of carbonyl compounds generated from the E-cigarette using coupled silica cartridges impregnated with hydroquinone and 2, 4-dinitrophenylhydrazine, followed by high-performance liquid chromatography. *Anal Sci.*, 29(12):1219–1222. doi:10.2116/analsci.29.1219.
- Wang, G.-W., Guo, Y., Vondriska, T. M., Zhang, J., Zhang, S., Tsai, L. L., ... Prabhu, S. D. (2008). Acrolein consumption exacerbates myocardial ischemic injury and blocks nitric oxide-induced PKC ϵ signaling and cardioprotection. *J Mol Cell Cardiol.*, 44(6):1016–1022. doi:10.1016/j.yjmcc.2008.03.020.